

On the Linearity of Free Nilpotent-by-Abelian Groups*

C. K. GUPTA AND N. D. GUPTA

Department of Mathematics, University of Manitoba, Winnipeg, Canada

Communicated by J. A. Green

Received August 24, 1971

INTRODUCTION

Linearity of free nilpotent groups seems to be well known. Among the other relatively free linear groups are: $F/\delta_2(F)$ (W. Magnus [6]), $F/\gamma_2\gamma_3(F) \cdot \gamma_3\gamma_2(F)$ (C. K. Gupta [3]) and $F_3/[\delta_2(F_3), F_3]$ (C. K. Gupta [2]). In this paper we show that $F/\gamma_n\gamma_2(F)$ ($n \geq 2$) is isomorphic to a linear group. This answers in the affirmative a recent question of B.A.F. Wehrfritz raised in Oberwolfach, May 1971.

NOTATION

Most of our notation will be developed as we proceed. However, the following notation is standard: $[a, b] = a^{-1}b^{-1}ab$; $[a, b, c] = [[a, b], c]$; $\gamma_m(A)$ is the m -th term of the lower central series of A ; $\delta_m(A)$ is the m -th term of the derived series of A ; $\gamma_r\gamma_s(A) = \gamma_r(\gamma_s(A))$. The free group of countable infinite rank or finite (unspecified) rank is denoted by F , whereas F_m denotes the free group of rank m .

PRELIMINARIES

Let F_m ($m \geq 2$) be the free group of rank m freely generated by a_1, \dots, a_m . Modulo $F_m' (= \gamma_2(F_m))$ define an order relation " $<$ " on the elements of F_m as follows:

$$u = a_1^{\alpha_1} \cdots a_m^{\alpha_m} < v = a_1^{\beta_1} \cdots a_m^{\beta_m} \quad (\alpha_i, \beta_j \in \mathbb{Z})$$

* Research supported in part by the National Research Council of Canada.

if

- (i) $\sum_{i=1}^m |\alpha_i| < \sum_{i=1}^m |\beta_i|$; or
- (ii) $\sum_{i=1}^m |\alpha_i| = \sum_{i=1}^m |\beta_i|$, and there is a least integer $t \in \{1, \dots, m\}$ such that $|\alpha_i| = |\beta_i|$ for $i < t$ and $|\alpha_t| < |\beta_t|$; or
- (iii) $|\alpha_i| = |\beta_i|$ for all $i \in \{1, \dots, m\}$ and there is a least integer $t \in \{1, \dots, m\}$ such that $\alpha_i = \beta_i$ for $i < t$ and $\alpha_t < \beta_t$.

For each pair (i, j) , $m \geq i > j \geq 1$, put

$$c_{ij}(k) = [a_i, a_j]^{u_{ij}(k)},$$

where $u_{ij}(k) = a_{jk}^{\alpha_1} \cdots a_{mk}^{\alpha_m}$ for some $\alpha_{jk}, \dots, \alpha_{mk} \in \mathbb{Z}$.

We extend the order relation " $<$ " to $c_{ij}(k)s'$ as follows: define

$$c_{ij}(k) < c_{i'j'}(k')$$

if

- (i) $u_{ij}(k) < u_{i'j'}(k')$; or
- (ii) $u_{ij}(k) = u_{i'j'}(k')$ and $i < i'$; or
- (iii) $u_{ij}(k) = u_{i'j'}(k')$, $i = i'$ and $j < j'$.

Let $C = \{c_{ij}(k), m \geq i > j \geq 1, k = 1, 2, \dots\}$. Using the above order relation we may re-arrange the contents of C as

$$B_1(=C) = \{b_{11}, b_{12}, \dots, | b_{1i} < b_{1j} \text{ for } i < j\}.$$

Using B_1 as an ordered set of symbols we let B_n denote the inductively defined and ordered set of (standard) basic commutators of weight n ($n \geq 1$), B_1 being the ordered set of basic commutators of weight 1 (see, for instance, M. Hall [5, Chapter 11]). Thus for each $n \geq 1$ we may write

$$B_n = \{b_{n1}, b_{n2}, \dots\}, \quad \text{where} \quad b_{n1} < b_{n2} < \dots.$$

We now prove,

LEMMA 1. *Modulo $\gamma_{n+1}(F_m')$, B_n generates $\gamma_n(F_m')$.*

Proof. Modulo $\gamma_2(F_m')$, F_m' is clearly generated by all elements of the form $[a_i, a_j]^w$ ($i > j$), where $w = a_1^{\alpha_1} \cdots a_m^{\alpha_m}$. Using the standard identities

$$[a_i^{\epsilon_i}, a_j^{\epsilon_j}] = [a_i, a_j]^{\epsilon_i \epsilon_j a_i^{\epsilon_i(\epsilon_i-1)/2} a_j^{\epsilon_j(\epsilon_j-1)/2}}, \quad [a, b, c] = [a, c, b][b, c, a]^{-1}$$

modulo $\gamma_2(F_m')$ and $[a, b]^c = [a, b][a, b, c](\epsilon_i, \epsilon_j = \pm 1)$, it follows that $[a_i, a_j]^{a_k^{\epsilon_k}}$ ($i > j > k$) $\in \text{gp}\{B_1\}$ modulo $\gamma_2(F_m')$. Thus by repeated application

of this last observation it follows that $[a_i, a_j]^w$ ($i > j$) $\in gp\{B_1\}$ for all $w \in F_m$. This proves the lemma for $n = 1$. Now modulo $\gamma_{n+1}(F_m')$, $\gamma_n(F_m')$ is generated by all left-normed commutators of weight n whose entries are the generators of F_m' , namely, b_{1k} 's. In other words, $\gamma_n(F_m')$ is generated by all commutators of the form

$$[b_{1i_1}, \dots, b_{1i_n}].$$

But $[b_{1i_1}, \dots, b_{1i_n}] \in gp\{B_n\}$ and we have the required proof.

Remark 1. As an application of our main result we shall also show that $\gamma_n(F_m')$ is freely generated by B_n modulo $\gamma_{n+1}(F_m')$ for $n \geq 1$ (c.f. Bachmuth [1] for the case $n = 1$).

Let H be the free abelian group of rank $2m$ ($m \geq 2$) freely generated by $y_1, \dots, y_m, z_1, \dots, z_m$ and let $P = \{\rho_1, \rho_2, \dots\}$ be a set of independent and commuting indeterminates. Consider $S = ZH[P]$ to be the polynomial ring in ρ 's over the integral group ring ZH . For each pair (i, j) , $m \geq i > j \geq 1$, define

$$\mu_{ij} = y_i^{-1} z_i^{-1} (-1 + y_j) \rho_i - y_j^{-1} z_j^{-1} (-1 + y_i) \rho_j.$$

We prove

LEMMA 2. In S , suppose $\sum_{(i,j)=(2,1)}^{(m,m-1)} (\sum_{k=1}^{t_{ij}} \zeta_{ij}(k) w_{ij}(k) \mu_{ij}) = 0$, where

(i) $\{\zeta_{ij}(k), k = 1, \dots, t_{ij}\}$ is a set of elements of S such that each term of $\zeta_{ij}(k)$ is of degree zero in each $y_1, \dots, y_m, \rho_1, \dots, \rho_m$; and

(ii) $\{w_{ij}(k), k = 1, \dots, t_{ij}\}$ is a set of words in y 's of the form $y_{i_1}^{\alpha_{i_1 k}} \cdots y_m^{\alpha_{m k}}$ with $w_{ij}(1) < \cdots < w_{ij}(t_{ij})$ (the ordering being parallel to that of F_m/F_m'). Then $\zeta_{ij}(k) = 0$ for all (i, j) ($m \geq i > j \geq 1$) and all $k \in \{1, \dots, t_{ij}\}$.

Proof. Put $\tau_{ij} = \sum_{k=1}^{t_{ij}} \zeta_{ij}(k) w_{ij}(k)$. Then we first of all prove by induction on $m \geq 2$, that $\tau_{ij} \mu_{ij} = 0$ for all (i, j) , $m \geq i > j \geq 1$. If $m = 2$, there is nothing to prove. Let $m > 2$ and assume that the result holds for $m - 1$. We have

$$\begin{aligned} 0 &= \sum_{(i,j)=(2,1)}^{(m,m-1)} \tau_{ij} \mu_{ij} \\ &= \sum_{j=1}^{m-1} \tau_{mj} \mu_{mj} + \sum_{(i,j)=(2,1)}^{(m-1,m-2)} \tau_{ij} \mu_{ij}. \end{aligned}$$

Thus by the induction hypothesis it is enough to show that $\tau_{mj} \mu_{mj} = 0$ for

$j = 1, \dots, m-1$. Since ρ 's are independent, the sum of the coefficients of ρ_m in $\sum_{(i,j)=(2,1)}^{(m,m-1)} \tau_{ij} \mu_{ij}$ must be zero. Accordingly, we have

$$\begin{aligned} 0 &= \sum_{j=1}^{m-1} \tau_{mj} (y_m^{-1} z_m^{-1} (-1 + y_j)) \\ &= (y_m^{-1} z_m^{-1}) \sum_{j=1}^{m-1} \tau_{mj} (-1 + y_j) \\ &= \sum_{j=1}^{m-1} \tau_{mj} (-1 + y_j) \\ &= \tau_{m1} (-1 + y_1) + \xi, \end{aligned}$$

where each term in $\xi = \sum_{j=2}^{m-1} \tau_{mj} (-1 + y_j)$ is of degree zero in y_1 (by definition). Writing

$$\tau_{m1} = \sum_{i=-t}^{+t} \xi_i y_1^i,$$

where ξ_i 's are all of degree zero in y_1 , we have

$$0 = \sum_{i=-t}^{+t} \xi_i y_1^i (-1 + y_1) + \xi.$$

Equating the coefficients of y_1^i to 0 for all $i = -t, \dots, t+1$, we get

$$\begin{aligned} 0 = \xi_{-t} &= \xi_{-t} - \xi_{-t+1} = \dots = \xi_{-2} - \xi_{-1} = \xi_{-1} - \xi_0 + \xi = \xi_0 - \xi_1 \\ &= \dots = \xi_{+t-1} - \xi_t = \xi_t. \end{aligned}$$

Solving all these equations we see that

$$0 = \xi_{-t} = \dots = \xi_{+t} = \xi,$$

and in particular $\tau_{m1} = 0$. Similarly $0 = \tau_{m2} = \dots = \tau_{m,m-1}$. Thus we have shown that $\tau_{ij} \mu_{ij} = 0$ for $m \geq i > j \geq 1$.

Next we consider.

$$\begin{aligned} 0 &= \tau_{ij} \mu_{ij} \\ &= \sum_{k=1}^{t_{ij}} \zeta_{ij}(k) w_{ij}(k) \mu_{ij}. \end{aligned}$$

Since the sum of the coefficients of ρ_i in $\tau_{ij}\mu_{ij}$ is zero, we have

$$\begin{aligned} 0 &= \left(\sum_{k=1}^{t_{ij}} \zeta_{ij}(k) w_{ij}(k) \right) (y_i^{-1} z_i^{-1} (-1 + y_j)) \\ &= \sum_{k=1}^{t_{ij}} \zeta_{ij}(k) w_{ij}(k), \text{ since } S \text{ is an integral domain.} \end{aligned}$$

But $\zeta_{ij}(k)$'s are all of degree zero in y 's and $w_{ij}(k)$'s are all distinct words in y 's. Thus $\sum_{k=1}^{t_{ij}} \zeta_{ij}(k) w_{ij}(k) = 0$ implies that $\zeta_{ij}(k) w_{ij}(k) = 0$ and hence $\zeta_{ij}(k) = 0$ for all $k = 1, \dots, t_{ij}$. This completes the proof of the lemma.

As an immediate corollary to Lemma 2 we record,

LEMMA 3. *The conclusion of Lemma 2 remains valid if μ_{ij} is replaced by $\mu_{ij}^* = y_i^{-1}(-1 + y_j z_j^{-1}) \rho_i - y_j^{-1}(-1 + y_i z_i^{-1}) \rho_j$ and $w_{ij}(k) = y_j^{\alpha_{jk}} \cdots y_m^{\alpha_{mk}}$ is replaced by $w_{ij}^*(k) = (y_j^{\alpha_{jk}} \cdots y_m^{\alpha_{mk}})(z_j^{-\alpha_{jk}} \cdots z_m^{-\alpha_{mk}})$.*

Proof. Since H is freely generated by $y_1, \dots, y_m, z_1, \dots, z_m$ it is also freely generated by $y_1 z_1^{-1}, \dots, y_m z_m^{-1}, z_1, \dots, z_m$. Further since μ_{ij}^* and $w_{ij}^*(k)$ are obtained from μ_{ij} and $w_{ij}(k)$, respectively, by replacing each y_i by $y_i z_i^{-1}$ and $\zeta_{ij}(k)$'s are unaffected, we have the required proof.

THE MAIN RESULT

Let G be the free abelian group freely generated by

$$\{x_{ij}, i = 1, \dots, n+1 (n \geq 1), j = 1, 2, \dots\}$$

and let $A = \{\lambda_{i,i-1}^{(k)}, n+1 \geq i \geq 2, k = 1, 2, \dots\}$ be a set of independent and commuting indeterminates. Consider the polynomial ring $R = ZG[A]$ over the group ring ZG . Let M_{n+1} denote the multiplicative group of $n+1 \times n+1$ matrices over R generated by $\langle a_1 \rangle, \langle a_2 \rangle, \dots$, where

$$\langle a_k \rangle = \begin{bmatrix} x_{1k} & & & & & \\ \lambda_{21}^{(k)} & x_{2k} & & & & \\ 0 & \lambda_{32}^{(k)} & x_{3k} & & & \\ \vdots & \vdots & \vdots & & & \\ \vdots & \vdots & \vdots & & & \\ 0 & \cdot & \cdot & 0 & \lambda_{n+1,n}^{(k)} x_{n+1,k} \end{bmatrix} \quad \bigcirc$$

for $k = 1, 2, \dots$. Let F be the free group freely generated by a_1, a_2, \dots . If $w = a_{i_1}^{\epsilon_1} \cdots a_{i_l}^{\epsilon_l}$ ($\epsilon_i \in \{-1, 1\}$) is an arbitrary word in F , we define

$$\langle w \rangle = \langle a_{i_1} \rangle^{\epsilon_1} \cdots \langle a_{i_l} \rangle^{\epsilon_l},$$

so that the mapping

$$\varphi: w \rightarrow \langle w \rangle$$

defines a homomorphism of F onto M_{n+1} . Our main result may now be stated as

THEOREM. *The kernel of φ is equal to $\gamma_{n+1}(\gamma_2(F))$.*

To simplify calculations we introduce certain mappings

$$\alpha_{ij}(n+1 \geq i \geq j \geq 1)$$

of F into R defined as:

$$\alpha_{ij}(w) = ij\text{-entry of the matrix } \langle w \rangle, \text{ for all words } w \in F.$$

Further, to each $w = a_{i_1}^{\epsilon_1} \cdots a_{i_l}^{\epsilon_l}$ in F we define $\mathbf{w}(j) = x_{j i_1}^{\epsilon_1} \cdots x_{j i_l}^{\epsilon_l} \in G$ for $j = 1, \dots, n+1$. In particular, $\mathbf{w}(j) = 1$ if and only if $w \in F' (= \gamma_2(F))$. We record some of the consequences of our definitions in the following lemma.

LEMMA 4. *Let u, v, w, \dots be words in F , then*

- (i) $\alpha_{ii}(u) = \mathbf{u}(i)$;
- (ii) $\alpha_{ij}(e) = 0$ for $i > j$ (e is the identity of F);
- (iii) $\alpha_{ij}(uv) = \sum_{k=j}^i \alpha_{ik}(u) \alpha_{kj}(v)$;
- (iv) $\alpha_{i, i-1}(a_k) = \lambda_{i, i-1}^{(k)}$;
- (v) $\alpha_{i, i-1}(u^{-1}) = -\mathbf{u}_{(i-1)}^{-1} \mathbf{u}_{(i)}^{-1} \alpha_{i, i-1}(u)$;
- (vi) $\alpha_{i, i-1}[u, v] = \mathbf{u}_{(i)}^{-1}(-1 + \mathbf{v}_{(i-1)} \mathbf{v}_{(i)}^{-1}) \alpha_{i, i-1}(u) \\ - \mathbf{v}_{(i)}^{-1}(-1 + \mathbf{u}_{(i-1)} \mathbf{u}_{(i)}^{-1}) \alpha_{i, i-1}(v)$;
- (vii) $\alpha_{i, i-1}[u, v]^w = \mathbf{w}_{(i-1)} \mathbf{w}_{(i)}^{-1} \alpha_{i, i-1}[u, v]$;
- (viii) if $u \in \gamma_r \gamma_2(F)$ then $\alpha_{ij}(u) = 0$ for all $r > i - j$;
- (ix) if $u \in \gamma_r \gamma_2(F)$, $v \in \gamma_s \gamma_2(F)$ and $r + s = i - j$, then
$$\alpha_{ij}[u, v] = \alpha_{i, i-r}(u) \alpha_{i-r, j}(v) - \alpha_{i, i-s}(v) \alpha_{i-s, j}(u)$$
- (x) if $u, v \in \gamma_r \gamma_2(F)$ and $r = i - j$, then
$$\alpha_{ij}(u^l) = l \alpha_{ij}(u) \quad \text{and} \quad \alpha_{ij}(uv) = \alpha_{ij}(u) + \alpha_{ij}(v).$$

Proof. The proofs of (i)–(iv) are trivial. For the proof of (v) note by (i)–(iv) that

$$\begin{aligned} 0 &= \alpha_{i,i-1}(e) = \alpha_{i,i-1}(uu^{-1}) \\ &= \alpha_{i,i-1}(u) \alpha_{i-1,i-1}(u^{-1}) + \alpha_{i,i}(u) \alpha_{i,i-1}(u^{-1}) \\ &= \mathbf{u}_{(i-1)}^{-1} \alpha_{i,i-1}(u) + \mathbf{u}_{(i)} \alpha_{i,i-1}(u^{-1}). \end{aligned}$$

The proofs of (vi) and (vii) can be derived by repeated applications of (iii) and (v). For the proof of (viii) we refer the reader to formula (6) of Gupta–Gupta [4]. Lemma 3(b) of [4] together with (viii) proves (ix). Finally (x) follows by the product rule (iii) and (viii).

For $i - j \in \{1, \dots, n\}$ let $(R_{i-j}, +)$ denote the additive subgroup of R generated by all products of the form

$$\alpha_{i,i-1}(b_{1k_1}) \cdots \alpha_{j+1,j}(b_{1k_{i-j}}),$$

where b_{1k} 's are from B_1 . By Lemma 4(vi), (vii) we can identify each $\alpha_{r,r-1}(b_{1k})$ with an element of the form $zw_{ij}^*(k) \mu_{ij}^*$ (as given in Lemma 3). Further we note that if $r \neq s$ then $\alpha_{r,r-1}(b_{1k})$ and $\alpha_{s,s-1}(b_{1k})$ involve independent symbols. Thus by repeated applications of Lemma 3 we have the following important lemma.

LEMMA 5. $(R_{i-j}, +)$ is freely generated by products of the form $\alpha_{i,i-1}(b_{1k_1}) \cdots \alpha_{j+1,j}(b_{1k_{i-j}})$.

Let N denote the free associative ring generated by v_1, v_2, \dots and let $(N_{i-j}, +)$, $i - j \in \{1, \dots, n\}$ denote the additive subgroup of N generated by all elements of the form

$$v_{k_1} \cdots v_{k_{i-j}}.$$

By Lemma 5, the mapping

$$\chi_{ij} : v_{k_1} \cdots v_{k_{i-j}} \rightarrow \alpha_{i,i-1}(b_{1k_1}) \cdots \alpha_{j+1,j}(b_{1k_{i-j}})$$

defines an isomorphism of $(N_{i-j}, +)$ onto $(R_{i-j}, +)$. By Lemma 4(ix), it is easy to see that if

$$\beta(b_{1k_1}, \dots, b_{1k_{i-j}}) \in B_{i-j}$$

for some bracket arrangement “ β ”, then

$$\alpha_{ij}(\beta(b_{1k_1}, \dots, b_{1k_{i-j}})) \chi_{ij}^{-1} = \beta(v_{k_1}, \dots, v_{k_{i-j}})$$

is a basic Lie-element in N with the same bracket arrangement " β ", in which each occurrence of b_{1k} has been replaced by v_k . Thus if

$$\sum_{k=1}^t r_k \alpha_{ij}(b_{i-j,k}) = 0,$$

then we have in turn

$$\begin{aligned} \left(\sum_{k=1}^t r_k \alpha_{ij}(b_{i-j,k}) \right) \chi_{ij}^{-1} &= 0; \\ \sum_{k=1}^t r_k \alpha_{ij}(\beta_k(b_{1k_1}, \dots, b_{1k_{i-j}})) \chi_{ij}^{-1} &= 0; \\ \sum_{k=1}^t r_k \beta_k(v_{k_1}, \dots, v_{k_{i-j}}) &= 0; \end{aligned}$$

which implies that $r_k = 0$ for $k = 1, \dots, t$ since basic Lie-elements in N are linearly independent (M. Hall [5, Chapter 11]). We record this observation as

LEMMA 6. $\sum_{k=1}^t r_k \alpha_{ij}(b_{i-j,k}) = 0$ implies $r_k = 0$ for each $k = 1, \dots, t$.

PROOF OF THE MAIN THEOREM

By Lemma 4(viii),

$$\alpha_{ij}(u) = 0 \quad \text{for all } u \in \gamma_{n+1}\gamma_2(F)$$

and all (i, j) , $n+1 \geq i > j \geq 1$. Thus $\gamma_{n+1}\gamma_2(F) \subseteq \ker \varphi$, the kernel of φ .

For the other inclusion we let $w = a_{i_1}^{e_1} \cdots a_{i_t}^{e_t} \in \ker \varphi$. We proceed to show that $w \in \gamma_{n+1}\gamma_2(F)$. We may assume that $\gamma_{n+1}\gamma_2(F) = \{e\}$ and conclude $w = e$. Since w involves only finitely many generating symbols of F , we may assume that $w \in F_m$ and $\gamma_{n+1}\gamma_2(F_m) = \{1\}$. Since $w \in \ker \varphi$, $\alpha_{ii}(w) = 1$ for all i which implies that $w(i) = 1$ and hence $w \in \gamma_2(F)$. To complete the proof of the theorem it is enough to show that if for some $s \in \{1, \dots, n\}$, $w \in \gamma_s\gamma_2(F_m)$ then $w \in \gamma_{s+1}\gamma_2(F_m)$. Indeed if $w \in \gamma_s\gamma_2(F_m)$, then modulo $\gamma_{s+1}\gamma_2(F_m)$,

$$w = b_{s_1}^{r_1} \cdots b_{s_t}^{r_t}, \text{ by Lemma 1.}$$

Since $w \in \ker \varphi$, $\alpha_{s+1,1}(w) = 0$ and by Lemma 4(x) this implies that

$$\begin{aligned} 0 &= \alpha_{s+1,1}(b_{s_1}^{r_1} \cdots b_{s_t}^{r_t}) \\ &= \sum_{i=1}^t r_i \alpha_{s+1,1}(b_{s_i}), \end{aligned}$$

and by Lemma 6, $r_1 = \cdots = r_t = 0$. Thus $w \in \gamma_{s+1}\gamma_2(F_m)$. This completes the proof of the main theorem.

CONCLUDING REMARKS

Remark 2. As promised in Remark 1, let us suppose that $b_{n_1}^{r_1} \cdots b_{n_t}^{r_t} = e$ modulo $\gamma_{n+1}\gamma_2(F_m)$. Then we have

$$\begin{aligned} 0 &= \alpha_{n+1,1}(b_{n_1}^{r_1} \cdots b_{n_t}^{r_t}) \\ &= \sum_{r=1}^t r_i \alpha_{n+1,1}(b_{n_i}) \quad (\text{by Lemma 4(x)}) \end{aligned}$$

and hence $r_1 = \cdots = r_t = 0$ (by Lemma 6). We record this as

COROLLARY 1. *Modulo $\gamma_{n+1}(\gamma_2(F_m))$, $\gamma_n\gamma_2(F_m)$ is freely generated by B_n .*

Remark 3. By Lemma 4(v) we have $\alpha_{i,i-1}(a_k^{-1}) = -x_{k,i-1}^{-1}x_{k,i}^{-1}\lambda_{i,i-1}^{(k)}$. Using this, the product rule 4(iii) and $0 = \alpha_{i,i-2}(a_k a_k^{-1})$ we can compute $\alpha_{i,i-2}(a_k^{-1})$ and similarly $\alpha_{ij}(a_k^{-1})$ for all $i-j \in \{1, \dots, n\}$. It follows that if $w = a_{i_1}^{\epsilon_1} \cdots a_{i_t}^{\epsilon_t}$ is an arbitrary word of F then $\langle w \rangle = \langle a_{i_1} \rangle^{\epsilon_1} \cdots \langle a_{i_t} \rangle^{\epsilon_t}$ can be effectively computed. Since $ZG[A]$ has a solvable word problem, it can be decided whether or not $\langle w \rangle$ is the identity matrix. Thus we have the following,

COROLLARY 2¹. *$F/\gamma_n\gamma_2(F)$ has solvable word problem.*

Remark 4. Let $A = \{\lambda_{i,i-1}^{(k)}, n+1 \geq i \geq 2, k = 1, 2, \dots\}$ be a set of independent and commuting indeterminates and let $Z[A]$ be the polynomial ring over Z . Using the theory of basic commutators it can be seen that the multiplicative group of $n+1 \times n+1$ matrices over $Z[A]$ generated by

$$\begin{pmatrix} 1 & & & & \\ \lambda_{21}^{(k)} & 1 & & & \\ 0 & \lambda_{32}^{(k)} & 1 & & \\ \vdots & & & \ddots & \\ 0 & \cdots & 0 & \lambda_{n+1,n}^{(k)} & 1 \end{pmatrix}, \quad k = 1, 2, \dots$$

¹ The referee has pointed out that every finitely generated linear group has solvable word problem (M. O. Rabin, unpublished).

is isomorphic to $F/\gamma_{n+1}(F)$. According to a suggestion of Professor P. Hall (communicated to the second author a few years ago by Dr. H. Heineken) the same proof can be carried over if $Z[A]$ is replaced by $Z[\lambda]$ and $\lambda_{i,i-1}^{(k)}$ is replaced by $\lambda^{2^{nk+(i-2)}}$.

REFERENCES

1. S. BACHMUTH, Automorphisms of free metabelian groups, *Trans. Amer. Math. Soc.* **118** (1965), 93-104.
2. C. K. GUPTA, A faithful matrix representation for certain centre-by-metabelian groups, *J. Austral. Math. Soc.* **10** (1969), 451-464.
3. C. K. GUPTA, On free groups of the variety $\mathfrak{N}_2 \wedge \mathfrak{N}_2\mathfrak{N}$, *Canad. Math. Bull.* **13** (1970), 443-446.
4. C. K. GUPTA AND N. D. GUPTA, Some groups of prime exponent, *J. Combinatorial Theory* **5** (1968), 397-407.
5. MARSHALL HALL, JR., "The Theory of Groups," MacMillan, New York, 1960.
6. WILHELM MAGNUS, On a theorem of Marshall Hall, *Ann. of Math.* **40** (1939), 764-768.